

Soil irrigation frequencies, compaction, air porosity and shear stress effects on soybean root development

Frecuencias de riego, compactación, porosidad aerífera y tensión cortante del suelo relacionadas con el desarrollo radical de soya

G. A. Hossne*, J. Méndez*, M. Trujillo*, F. Parra*

ABSTRACT

Effect of soil compaction on the root development of soybean plants (*Glycine max* L. Merrill cv. San Baiba), subjected to different treatments of irrigation frequencies, air porosity and shear stress of a sandy loam soil were performed. Soybeans are important agricultural crop of the area, where the soil is incompressible and easy deformable. The objectives were to determine: (a) Consequences of water content, compaction and air porosity over the root length, root volume, and root fresh mass; and (b) Results of air content, shear stress and normal strain on root growth. The methods used were the Proctor test, triaxial, water meters, watering frequency and 30 cm × 30 cm × 1.5 cm plastic cylinders. The randomized block was used in simple factorial arrangement with four levels of compaction per layer (0, 12, 24, 36) and four soil water content through four irrigation frequencies (daily, inter-day, every two days and each three days). The findings were: (a) Root length between 24 cm and 79.5 cm; (b) Root volume from 2 cm³ to 40 cm³; (c) Root fresh mass between 7.58 g and 34.04 g, with a higher values tendency for daily and inter-day irrigation frequencies. The daily and inter day irrigation frequencies average results were above the grand mean (52.31 cm) of root length for the four levels of compaction. It was concluded that the soybeans root system was positively influenced by water content, more than compaction and the other variables under study.

RESUMEN

Se estudió el efecto de la compactación del suelo en el desarrollo de la raíz de plantas de soya (*Glycine max* L. Merrill cv. San Baiba) sometidas a diferentes tratamientos de frecuencias de riego, porosidad aerífera y tensión cortante de un suelo franco arenoso de sabana. La soya es de importancia agrícola en la zona, donde el suelo es incompresible y de fácil deformación. Los objetivos fueron determinar (a) el efecto del contenido de humedad y porosidad aerífera sobre la longitud, el volumen radical y la masa fresca radical, y (b) la relación del contenido aerífero, tensión cortante y tensión normal con el crecimiento radical. Se utilizó el método Proctor, triaxial, medidores de humedad, frecuencia de riego y cilindros plásticos de 30 cm × 30 cm × 1.5 cm. Se emplearon bloques al azar en arreglo factorial simple con cuatro niveles de compactación por capa (0, 12, 24, 36) y cuatro de humedad a través de cuatro frecuencias de riego (todos los días, interdiario, cada dos días y cada tres días). Entre los resultados: (a) la longitud radical entre 24 cm y 79.5 cm; (b) el volumen radical entre 2 cm³ y 40 cm³; (c) la masa fresca radical entre 7.58 g y 34.04 g, con tendencia de los valores mayores para los tratamientos con frecuencias de riego diaria e interdiaria que produjeron resultados promedios por encima de la gran media (52.31 cm) de la longitud radical para los efectos de los cuatro niveles de compactación. Se concluyó que el sistema radical fue más positivamente influenciado por la humedad que por los efectos de las otras variables estudiadas.

INTRODUCTION

Compaction approaches soil particles, dry or wet, reducing the air space and the soil ability to adequately preserve enough amounts of air. The soil pore space needed to ensure the gas exchange required by a healthy root system should be between 10% and 20% of air. According to Kozlowski (1985) and Costello, MacDonald & Jacobs (1991), the roots work best with a level above

Recibido: 8 de septiembre de 2014

Aceptado: 7 de noviembre de 2014

Keywords:

Savanna soil; compaction; irrigation; morphological characters.

Palabras clave:

Suelos de sabana; compactación; riego; caracteres morfológicos.

Cómo citar:

Hossne, G. A., Méndez, J., Trujillo, M. & Parra, F. (2015). Soil irrigation frequencies, compaction, air porosity and shear stress effects on soybean root development. *Acta Universitaria*, 25(1), 22-30. doi: 10.15174/au.2015.676

* Departamento de Ingeniería Agrícola, Núcleo de Monagas, Universidad de Oriente. Apartado postal núm. 414, Maturín, Estado Monagas, Venezuela. Correos electrónicos: americohossne-garcia@gmail.com; americohossne@cantv.net

10% oxygen, levels below 4%-5% highly inhibit root growth. The soil structure is changed by compaction, reducing size and pore continuity, causing increased soil density. It is conceived that the soil compaction, usually measured by evaluation of the dry density, creates root growth inhibition of plants, such as in soybeans which is agriculturally important for the area. It has been considered that compaction increases the soil resistance. According to Ponder (2004), the mechanism that compacted soil supported better growth more than not compacted is not completely understood. Much of the better growth is likely owing to better soil physical changes that caused better soil moisture conditions for growth. Severely compacted layer below the depth of tillage is a special problem of soil compaction in many farmlands. This severely compacted layer restricts root growth and water available for plant roots. Taylor & Burnett (1964) and Taylor (1974) reported on the excessive strength of plow pans when were dry; dryness was the main restriction hindering root growth, due that soil resistance diminished with rain or irrigation. The soil resistance critical value, at which plant roots do not elongate, is produced by soil moisture and stages of plant development. Comparison of the measures of compaction in the literature is quite difficult because of a lack of information on moisture and development status.

Compaction, by definition, is the reduction of air-filled pores of the soil. It is common to provide the soil bulk density measurements unreferenced with the values of moisture content of the soil. Miller, Scott & Hazard (1996), and Cochran & Brock (1985) reported existing ample evidence supporting the reduction in growth caused by soil compaction, it has not always produced negative growth effects. Gomez, Powers, Singer & Horwath (2002) specified that the effect of compaction varied with the texture and soil moisture. These authors also reported that compaction reduced the growth of young ponderosa pine in California on fine textured soils attributable to increased soil strength, but soil compaction increased growth on a sandy textured soil owing to increased water holding capacity. Rahman, Hara & Hoque (2005) conducted a greenhouse experiment, with sandy loam andisols soil, to evaluate the cause of different levels of compaction on soybean. Root dry weight, root shoot ratio and shoot dry weight decreased significantly with the increased energy levels and decreased with increased levels of compaction. Hossne *et al.* (2003); Hossne (2008) showed that soil bulk densities varied inversely with

soil water content and that soil shear stress was an inverse function of soil moisture, close to zero as it approached the soil liquid limit. Hossne (2004) reported that the action of root growth increased its axial stress for soil shear stress with a critical value of 2343.2 kPa for 20.78% soil moisture content, these soils exerted a shear strength is lower 500 kPa within the range of field capacity. Smucker & Erickson (1987) informed that plants subject to soil compaction are more susceptible to water stress. And according to Schumacher & Smucker (1984), root development may be damaged by lack of oxygen in compacted soils. The objectives of this study were to evaluate: (a) The influence of moisture content and air porosity levels of a sandy loam soil of savanna on the length, volume and fresh root mass of soybean; and (b) the relationship among soybean root growth and air porosity, shear strength and normal stress of the studied soil.

MATERIALS AND METHODS

A greenhouse cultivation of Soybeans (*Glycine max* L. Merrill cv. San Baiba) in sandy loam soil of savanna of the Monagas state, with the characteristics shown in table 1, was used in the study.

Table 1.
Soil physical characteristics and organic matter content of the cultivated soil.

Components	Size	Horizons
	mm	0 cm - 300 cm
		%
Very coarse sand	1	1.9055
Coarse sand	0.5	11.99
Medium sand	0.098	24.092
Fine sand	0.053	24.284
Very fine sand	0.041	9.303
Total sand		71.539
Silt		15.785
Clay (Kaolinite)		12.676
Organic matter		1.041
Textural class		SL

SL: United States Department of Agriculture (USDA) classification: SaLo (sandy loam).

Source: Authors own elaboration.

The Proctor method was used to determine the dry bulk density. The Proctor test is a geotechnical engineering test to find out the maximum density that may be practically achieved for a soil or similar substance. The Proctor soil compaction test is done by measuring the density, or dry unit weight of the soil being tested at different moisture content points. The aim of the soil test is usually to determine the optimum moisture content for the soil. In addition to soils, other substances, such as aggregate, gravel, or sand, may be measured. In this scenario, compacting the soil means increasing its density by forcing air out of the soil. By compacting the soil or aggregate at different moisture contents, an engineer may determine what the optimum moisture content and compaction level are of the soil or aggregate for a specific use in a particular engineering project. The Proctor test was invented in 1933 by Ralph R. Proctor (1933). Both the original and the modified Proctor tests also allow the use of a larger mold for measuring substances that contain larger particles, such as gravel. The air porosity (E_a) was determined using Equation 1, as a function of the total porosity (n), the volumetric water content (θ), dry bulk density (ρ_S), particle density (ρ_P), gravimetric water content (w) and the water density (ρ_W). Moisture soil meters and frequencies of irrigation were employed. The interaction of four compaction levels (Proctor strokes) per layer: (0, 12, 24, 36) and four soil water content through irrigation frequencies: (daily, inter-day, every two days and every three days) were studied. For determination of soil strength, $\tau = C + \sigma^* \tan(\phi)$, the triaxial apparatus was utilized. Sixty-four polymers cylinders 30 cm diameter, 30 cm height and 1.5 cm thick were employed, cut longitudinally into two halves at the center and fastened shown in figure 1.

$$E_a = n - \theta = \left(1 - \frac{\rho_S}{\rho_P}\right) - \frac{w \cdot \rho_S}{\rho_W} \quad (1)$$



Figure 1. Experiment view of the sixty-four treatments. Cylinders, Proctor hammer, Proctor bump support and moisture meters used in the study. Source: Authors own elaboration.

The harvest took place between seventy-two (72) and seventy-five (75) days. 24.57 kg (8.19 kg/layer) of soil per cylinder were utilized. Fertilizer application of 2.74 g/cylinder of 25 kg/ha-100 kg/ha-120 kg/ha of NPK was performed. One liter of water per irrigation frequency was applied. A randomized blocks experimental design (4*4) with simple factorial with sixteen (16) treatments (T_o) with four replications for a total of sixty-four (64) experimental units was applied. Irrigation were established: every day (Fr1), inter-day (Fr2), every two days (Fr3) and every three days (Fr4) and compaction levels: 0 drops/layer (C1), 12 drops/layer (C2), 24 drops/layer (C3) and 36 drops/layer (C4). Statistically regression analysis of variance (ANOVA), the least significant difference (LSD), the best subset regression method, and with multiple regressions it was introduced a cubic polynomial with the dependent variable root length (L) and the independent variables: compaction, represented by air porosity (E_a) and irrigation frequencies (Fr) were used.

RESULTS

By applying multiple regression, a third degree polynomial with thirteen terms was optimized, of which the terms Fr, $E_a \cdot Fr$, $E_a^2 \cdot Fr$, $E_a^2 \cdot Fr^2$, $E_a^3 \cdot Fr$, $E_a^3 \cdot Fr^2$ and $E_a^3 \cdot Fr^3$ were eliminated with p values greater than 0.05 in the ANOVA, to forge a third degree polynomial with six terms shown in equation 2, with R^2 of 92.33, adjusted R^2 of 87.21, standard error of 2.83 and absolute minimum error 1.56. The ANOVA p value of 0.0002 showed a statistically significant relationship between variables in a high level of significance. Figure 2 shows the response surface plot for the model of equation 2.

$$L = -262z69,7 + 1834,11 \cdot E_a - 42,622 \cdot E_a^2 + 17,163 \cdot Fr^2 + 0,33 \cdot E_a^3 - 0,011 \cdot E_a^2 \cdot Fr^3 + 0,00042 \cdot E_a^2 \cdot Fr \quad (2)$$




Figure 2. Root length response surface versus irrigation period and air porosity. Source: Authors own elaboration.

Figure 3 shows the results of the root length, shear stress and air porosity versus treatments and soil moisture recorded during the process of soybean root growth. Figure 4 evidence the results of root length, porosity and moisture versus the experiment treatments. Figure 5 indicates the results of radical fresh weight, air porosity and moisture versus the treatments. Figure 6 graphically displays the results of the rooting volume, air porosity and moisture versus treatments.

Figure 7 presents the root volume obtained with the treatments C4Fr4 (36 drops/layer with watering every three days) and C4Fr2 (36 drops/layer and inter-day watering). This supports the results presented in graphs and statistical analysis. According to Fageria, Balingar & Clark (2006), Grzesiak, Hura, Grzesiak & Pienkowski (1999), Iijima & Kono (1991), Masle (2002) and Yamauchi (1993) the inhibition of plant growth is attributed mainly to reduced rooting volume.




Figure 3. Root length versus treatments, air porosity and shear stress. Source: Authors own elaboration.




Figure 4. Root length (L) versus treatments, air porosity (Ea) and soil humidity (w). Source: Authors own elaboration.




Figure 5. Root fresh mass (M_{FR}) versus treatments, air porosity (Ea) and soil moisture. Source: Authors own elaboration.




Figure 6. Rooting volume (R_{SV}) versus soil treatments, soil air porosity and soil moisture. Source: Authors own elaboration.




Figure 7. C4Fr4 and C4Fr2 treatments view at the end of the experiment. Source: Authors own elaboration.

Table 2 presents the statistical results obtained from the least significant difference analysis and analysis of variance.

DISCUSSION

Root length versus irrigation period and air porosity, shown in response surface of figure 2, demonstrates that soil moisture was the variable almost influential for plant root development. The air porosity, with accepted percentage product of soil compacting effect, did not inhibited root growth, but fewer prevalent than water application as it is the intense blue shadow path seen adjacent to the irrigation frequency coordinate that exhibits the greatest root length assessments as noted in the palette values.

Table 2.

Mean analysis by least significant difference and analysis of variance (ANOVA) for root length (L) (cm) for four levels of compaction (C) and four irrigation frequencies (Fr) that formed the treatments for a savanna sandy loam soil of the Monagas state in Venezuela.

Independent variables		Dependent variable			
		Root length (L)			
Group 1	Group 2	Mean 1	Group 1	Mean 2	Group 2
C1Fr1	C4Fr3	65.08	A	50.64	BCDEF
C4Fr1	C4Fr3	60.98	AB	49.80	CDEF
C1Fr2	C2Fr3	60.81	ABC	49.71	DEF
C4Fr2	C2Fr4	59.09	ABCD	48.19	DEFG
C3Fr1	C3Fr4	56.00	ABCDE	47.35	EFG
C2Fr2	C3Fr2	55.96	ABCDE	46.75	EFG
C2Fr1	C4Fr4	54.84	ABCDE	41.90	FG
C1Fr4	C1Fr3	51.86	BCDEF	38.07	G

Least significant difference (LSD). Pairwise comparisons ($p \leq 0.05$). Different letters indicate statistically different averages. Critical value of 1.98 T and the critical value for the comparison were of 11.02.

Variance analysis (ANOVA) for root length growth (L)

Sources	DF	Sum of square	Mean square	F	P
Treatment	15	6313.8	420.92	3.40	0.0001
Error	112	13 852.2	123.68		
Total	127	20 166.0			
Grand mean			52.31		
VC			21.26		

$F_{Critical} 0.001(15:112) = 2.803$ y $F_{Critical} 0.05(15:112) = 1.756$. As $3.40 > 2.803$ and $3.40 > 1.76$, the null hypothesis was rejected, concluding that the results of the treatments were different.

Source: Authors own elaboration.

Figure 3, that presents the curves of shear stress (τ) for different chamber pressures, shows that the resistance of the studied soil under satisfactory soil moisture conditions (Fr1 and Fr2), became imperceptible. In this analysis, the graphs for the results of fresh mass and root volume are not shown attributable to the same trends obtained. It is seen that soil moisture was the almost influential upon root growth. At the highest levels of resistance of the soil, the roots are more sensitive to water deficit (Davis, 1984).

Figures 3, 4, 5, 6 and 7, show that the best root expansion of root length, fresh mass and root volume were for every day and inter-day irrigation periods, increasing with soil moisture increments. This coincides with the statistical results already presented.

It is observed in table 2 that the longest root length was achieved with the daily watering frequency for compaction (C1) and maximum compaction (C4). There were seven treatment groups without significant difference; this was expected, since the four levels of compaction (C1, C2, C3 and C4) did not produce an effect on the two soil water content levels: inter-day and daily irrigation. Also, the use of both frequencies were indifferent one another. This indicates that moisture was the parameter that most influenced the root growth. Thereon; Ehlers, Köpfe, Hesse & Böhm (1983) reported that mechanical impediments increased as dry bulk density increased and decreased by amplifying the water content. Similar results were reported by Blouin, Schmidt, Bulmer & Krzic (2004) and Coile (1948). According to Blackwell, Ward, Lefevre & Cowan (1985), and Boone, De Smet & van Loon (1985), soil wetness may produce an effect the rootability of pores smaller than the root diameter. It was observed that in compacted soil some roots were able to widen pore spaces by deforming the space with root expansion. Such deformation is easier for roots when soil is wet (low soil strength) than when it is dry (high soil strength). This was notified by Hossne (2004).

By using the best subsets regression analysis applied through the origin with independent variables moisture (w), dry bulk density (ρS) and treatment (To) as forced independent variables, and soil resistance (τ) and air porosity (Ea) as not enforced; the best models, in the order of influence, for root length (L) were w , ρS , To (R^2 0.946) and w ρS To Ea (R^2 0.949). This strengthens that soil water content was the crucial component that influenced root growth.

According to Smucker & Erickson (1987) plants subject to compaction are more susceptible to water

stress. Schumacher & Smucker (1984) specified that root development may be damaged by lack of oxygen in compacted soils. The lack of air porosity may cause harmful effects on plant growth (Drew, 1990; Grichko & Glick, 2001; He, Finlayson, Drew, Jordan & Morgan, 1996; Voorhees, Farrel & Larson, 1975 and Zou, Penfold, Sands, Misra & Hudson, 2001). When the proportion of oxygen diffusion decreases to 10% or less, it causes root injuries disabling its ability to function. Engelaar & Yoneyama (2000) and Beutler, Centurion & Da Silva (2005), found that the air porosity tightening was much larger than the field capacity, thus causing little restriction on soybean root development during the period; also, they found that the smallest value of penetrometer resistance caused reduction in soybean production was probably attributable to the large variation of soil water content during the cycle of soybeans. In this work the air porosity was much above 10%. These results are supported by Blouin *et al.* (2004); Hossne *et al.* (2003); Hossne (2004).

Soybean roots appear to grow in soil under conditions of low water stress (Boyer, Johnson & Saupe, 1980; Garay & Wilhelmy 1983; Sivakumar, Taylor & Shaw, 1977). Taylor, Roberson & Parker (1967) established relationships between soil strength, water content, the emergence of seedlings and root growth. These studies indicated that the growth of plants and roots were reduced when soil resistance reached critical levels consequential to natural or provoked compaction, concluding that the resistance should be considered as a main factor of impact. In this experiment for the studied soil, strength (τ) in humid conditions near field capacity was not influential. Compaction may affect the rate of elongation of roots. Taylor & Ratliff (1969) showed that an increased penetration resistance of 1.9 MPa decreased the speed of peanut root elongation to 50% of the maximum rate (2.7 mm per hour) through loose loamy sand. The same reduction was noted for cotton root with only an increased resistance of 0.72 MPa. Using spring wheat, Collis-George & Yoganathan (1985) showed that increasing shear strength of a fine sand seed bed from 19 kPa to 52 kPa reduced seminal root elongation from 43.5 cm to 0.2 cm per day. The limiting root strengths at low densities were achieved when the soil remained in a state of dryness. At 2.96 MPa no penetration occurred, caused by low water content or apparent density. According to Carter (1989) and Salter & Goode (1967) water stress was the most important soil factor limiting agricultural production in the world.

Savanna sandy soils large pores contain kaolinitic clays that are considered inexpandible but suitable

farm machine processes. In this regard, Goldsmith, Silva & Fischenich (2001) reported that sandy soils have large continuous pore, clays have small pores that transmit water slowly. Clays, however, contain more pore space than sandy soils. For growing plants, pore sizes are more important than total pore space. Therefore, plants will have a better environment in sandy soils if porosity is reduced because of the increase in water retention. The converse is true for clays. High porosity clays have a high macro movement that provides high infiltration and more water available for plants. Consequently, the plants will have a better environment in sandy soils if the porosity is decreased, then increasing moisture retention. According to Kim (2000) roots survive and grow where the water needed is available, temperatures are warm, and oxygen is present. The ability of primary root tips to enter, open and extend through the pores of the soil depends on the force generated by the root and penetration resistance. The atmosphere of the soil should contain an air space between 12% and 60%, oxygen content between 3% and 21% for root growth. Tokunaga (2006) concluded in his study that biomass production was greatest when water was readily available. In field work, where water availability may be very variable, when the compaction affects the availability of water may be more important than physical impediments. Using cotton planted in a fine sandy loam, Taylor & Gardner (1963) found that soil strength was critical in stopping root growth. They did not find a critical root limiting bulk density because root penetration at each bulk density was affected by soil water content. Root-limiting resistances were reached at lower bulk densities when soils were dry. Higher strength decreased root growth rate. No penetration occurred at 2.96 MPa regardless caused by low water content or bulk density.

CONCLUSIONS

The results were: (a) The root length varied between 24 cm and 79.5 cm; (b) The root volume between 2 cm³ and 40 cm³; (c) The radical fresh mass between 7.58 g and 34.04 g; (d) The soil rooted volume between of 83.0 cm³ 99 cm³, with a tendency of higher values for treatments with daily and inter-day irrigation frequencies. For daily and inter-day irrigation frequencies, root length mean scores were above the grand mean (52.31 cm) for four levels of compaction.

Root development was largely influenced by the soil moisture content. The consequence of the compaction attributable to the volume change caused by the Proc-

tor hammer drops showed their influence, but possibly owing to reduced air availability and not favored by soil compaction. Soil resistance, that is an inverse function of the moisture, had no influence when applied irrigation frequencies that produced the greatest soil moisture.

The root volume, root fresh weight and soil rooted volume conserved the tendency of higher values for treatments with daily and inter-day irrigation frequencies.

ACKNOWLEDGMENTS

The authors is grateful to the Research Council of the Universidad de Oriente in Venezuela for its support and funding for this research.

REFERENCES

- Beutler, A. N., J. F. Centurion & Da Silva, A. Pi (2005). Soil Resistance to Penetration and Least Limiting Water Range for Soybean Yield in a Haplustox from Brazil. *Brazilian Archives of Biology and Technology Journal*, 48(6), 863-871.
- Blackwell, P. S., Ward, M. A., Lefevre, R. N. & Cowan, D. J. (1985). Compaction of a swelling clay soil by agricultural traffic, effects upon conditions for growth of winter cereals and evidence for some recovery of structure. *Journal Soil Science*, 36, 633-636.
- Blouin, V., Schmidt, M., Bulmer, C. & Krzic, C. (december, 2004). Soil compaction and water content effects on lodgepole pine seedling growth in British Columbia. *SuperSoil 2004: 3rd Australian New Zealand Soils Conference, University of Sydney, Australia*. Published on CDROM. Retrieved January 15, 2014, from www.regional.org.au/au/asssi/
- Boone, F. R., De Smet, L. A. H. & van Loon, C. D. (1985). The effect of ploughpan in marine loam soils on potato growth I. Physical properties and rooting patterns. *Potato Research*, 28, 295-314.
- Boyer, J. S., Johnson, R. R. & Saupe, S. G. (1980). Afternoon water deficits and grain yields in old and new soybean cultivars. *Agronomy Journal*, 72, 981-982.
- Carter, T. E. (march, 1989). Breeding for drought tolerance in soybean: where do we stand? World Research Conference IV. Asociación Argentina De La Soja, Buenos Aires, Argentina.
- Cochran, P. H. & Brock, T. (1985). *Soil compaction and initial height growth of planted ponderosa pine*. Res. Note PNW-434. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station.
- Coile, T. S. (1948). Relation of soil characteristics to site index of loblolly and shortleaf pines in the Lower Piedmont region of North Carolina. *Duke University School of Forestry Bulletin*, 13, 78.
- Collis-George, N. & Yoganathan, P. (1985). The effect of soil strength on germination and emergence of wheat (*Triticum aestivum* L.) I. Low shear strength conditions. *Australian Journal of Soil Research*, 23, 577-587.

- Costello, L. R., MacDonald, J. D. & Jacobs, K. A. (1991). Soil Aeration and tree health: correlating soil oxygen measurements with the decline of established oaks. *USDA Forest Service General Technical Report PSW-126*, 295-299.
- Davis, G. R. (1984). Effect of soil compaction on root growth of *Pinus radiata* D. Don. *Proceeding Symposium Site and Productivity of Fast Growing Plantations*, 2, 871-877.
- Drew, M. C. (1990). Sensing soil oxygen. *Plant, Cell, and Environment*, 13, 681-693.
- Ehlers, W., Köpfe, U. Hesse, F. & Böhm, W. (1983). Penetration resistance and root growth of oats in tilled and untilled loess soils. *Soil & Tillage Research*, 3, 261-275.
- Engelaar, W. M. H. G. & Yoneyama, T. (2000). Combined effects of soil waterlogging and compaction on rice (*Oryza sativa* L.) growth, soil aeration, soil N transformations and N discrimination. *Biology and Fertility of Soils*, 32, 484-493.
- Fageria, N. K., Balingar, V. C. & Clark, R. B. (2006). *Physiology of crop production*. New York, London, Oxford: The Haworth Press Inc.
- Garay, A. F. & Wilhelmy, W. (1983). *Root System Characteristics of Two Soybean Isolines Undergoing Water Stress Condition*. Nebraska: USDA Agricultural Research Service. USDA-ARS / UNL Faculty. University of Nebraska. Retrieved January 15, 2014, from <http://digitalcommons.unl.edu/usdaarsfacpub>.
- Goldsmith, W., Silva, M. & Fischenich, C. (2001). *Determining optimal degree of soil compaction for balancing mechanical stability and plant growth capacity*. ERDC TN-EMRRP-SR-26. U.S. Army Engineer Research and Development Center, Vicksburg, MS.
- Gomez, A., Powers, R. F., Singer, M. J. & Horwath, W.R. (2002). Soil compaction effects on growth of young ponderosa pine following litter removal in California's Sierra Nevada. *Soil Science Society of America*, 66, 1334-1343.
- Grichko, V. P. & Glick, B. R. (2001). Ethylene and flooding stress in plants. *Plant Physiology*, 39, 1-9.
- Grzesiak, S., Hura, T., Grzesiak, M. T. & Pieńkowski, S. (1999). The impact of limited soil moisture and waterlogging stress conditions on morphological and anatomical root traits in maize (*Zea mays* L.) hybrids of different drought tolerance. *Acta Physiologiae Plantarum*, 21, 305-315.
- He, C., Finlayson, S. A., Drew, M. C., Jordan, W. R. & Morgan, P. W. (1996). Ethylene biosynthesis during aerenchyma formation in roots of maize subjected to mechanical impedance and hypoxia. *Plant Physiology*, 112, 1679-1685.
- Hossne, G. A. (2004). Evaluación terramecánica del crecimiento radical en un suelo Ultisol de sabana del estado Monagas, Venezuela. *UDO Agrícola*, 4(1), 45-52.
- Hossne, G. A. (2008). La densidad aparente y sus implicaciones agrícolas en el proceso expansión/contracción del suelo. *Terra Latinoamericana*, 26, 195-202.
- Hossne, G. A., Christopher, J., Paredes, G., Carvajal, E., Santaella, E., Malaver, J. & Cova, A. (2003). Resistencia terramecánica en función de la humedad edáfica de un suelo ultisol de sabana del estado Monagas, Venezuela. *Bio-agro*, 15(3), 173-181.
- Iijima, M. & Kono, Y. (1991). Inter specific differences of the root system structures of four cereal species as affected by soil compaction. *Japanese Journal Crop Science*, 60, 130-138.
- Kim, K. D. (2000). *The root growth requirements*. EUA: University of Georgia Warnell School of Forest Resources. Extension Publication FOR00-5. Retrieved November 25, 2013, from www.forestry.uga.edu/efr
- Kozłowski, T. T. (1985). Soil aeration, flooding and tree growth. *Journal of Arboriculture*, 11, 85-96.
- Masle, J. (2002). High soil strength: mechanical forces at play on root morphogenesis and in root:shoot signaling. In Y. Waisel, A. Eshel & U. Kafkafi U (Eds.), *Plant Roots the Hidden Half* (pp. 807-819). New York / Basel: Marcel Dekker Inc.
- Miller, R. E., Scott, W. & Hazard, J. W. (1996). Soil compaction and conifer growth after tractor yarding at three coastal Washington locations. *Canadian Journal of Forest Research*, 26, 225-236.
- Ponder, F. (2004). Soil compaction affects growth of young shortleaf pine following litter removal and weed control in the Missouri Ozarks. *Proceedings of the 14th Central Hardwoods Forest Conference* (pp. 255-264).
- Proctor, R. (1933). Fundamental principles of soil compaction. *Engineering News Record*, 111(9), 245-248.
- Rahman, M. H., Hara, M. & Hoque, S. (2005). Growth and nutrient uptake of grain legumes as affected by induced compaction in andisols. *International Journal of Agriculture and Biology*, 7, 740-743.
- Salter, P. J. & Goode, J. E. (1967). Crop response to water at different stage of growth. *Commonwealth Agricultural Bureaux*. Farnham Royal, Bucks England, 246.
- Schumacher, T. E. & Smucker, A. J. M. (1984). Effect of soil compaction on root growth and uptake of phosphorus. *Plant and Soil*, 77, 15-28.
- Sivakumar, M. V. K., Taylor, H. M. & Shaw, R. H. (1977). Top and root relations of field grown soybeans. *Agronomy Journal*, 69, 470-473.
- Smucker, A. J. M. & Erickson, A. E. (1987). Anaerobic stimulation of root exudates and diseases of peas. *Plant Soil*, 99, 423-433.
- Taylor, H. M. (1974). Root behavior as affected by soil structure and strength. In *The Plant Root and Its Environment* (pp. 271-291). Carson, E.W.: Ed. Univ. Press of Virginia, Charlottesville.
- Taylor, H. M. & Burnett, E. (1964). Influence of soil strength on root-growth habits of plants. *Soil Science*, 98, 174-180.
- Taylor, H. M. & Gardner, H. R. (1963). Penetration of cotton seedling taproots as influenced by bulk density, moisture content, and strength of soil. *Soil Science*, 96, 153-156.
- Taylor, H. M. & Ratliff, L. F. (1969). Root elongation rates of cotton and peanuts as a function of soil strength and soil water content. *Soil Science*, 108, 113-115.
- Taylor, H. M., Roberson, G. M. & Parker, J. J. (1967). Cotton seedling taproot elongation as affected by soil strength changes induced by slurring and water extraction. *Soil Science Society of American Proceedings*, 31, 700-704.

- Tokunaga, A. A. (2006). Effects of bulk density and soil strength on the growth of Blue wildrye (*Elymus glaucus* buckl) (Thesis of master). Humboldt State University, USA.
- Voorhees, D., Farrel, D. A. & Larson, W. E. (1975). Soil strength and aeration effects on root elongation. *Soil Science Society of America Journal*, 39, 948-953.
- Yamauchi, A. (1993). Significance of root system structure in relation to stress tolerance in cereal crop. In *Low-input sustainable crop production system in Asia* (pp. 347-360). Korea: Ksorean Society for Crop Science.
- Zou, C., Penfold, C., Sands, R., Misra, R. K. & Hudson, I. (2001). Effects of soil air-filled porosity, matric potential and soil strength on primary root growth of radiata pine seedlings. *Plant and Soil*, 236, 105-115.